

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G01N 33/68, 33/569, C07K 14/47, 16/18	A1	(11) International Publication Number: WO 00/22438 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23115 (22) International Filing Date: 5 October 1999 (05.10.99) (30) Priority Data: 09/169,574 9 October 1998 (09.10.98) US (71) Applicant (for all designated States except US): THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th floor, Oakland, CA 94607-5200 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PRUSINER, Stanley, B. [US/US]; 400 Pacheco Street, San Francisco, CA 94116 (US). SAFAR, Jiri, G. [CZ/US]; 1003 Stimel Drive, Concord, CA 94518 (US). (74) Agent: BOZICEVIC, Karl; Bozicevic, Field & Francis LLP, Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ASSAY FOR DISEASE RELATED CONFORMATION OF A PROTEIN		
(57) Abstract <p>An assay method is disclosed which isolates and detects the presence of a disease related conformation of a protein (e.g., PrP^{Sc}) present in a sample also containing the non-disease related conformation of the protein (e.g., PrP^C). The sample is treated (e.g., contacted with protease) in a manner which hydrolyzes the disease related conformation and not the non-disease related conformation. The treated sample is contacted with a binding partner (e.g., a labeled antibody which binds PrP^{Sc}) and the occurrence of binding provides an indication that PrP^{Sc} is present. Alternatively the PrP^{Sc} of the treated sample is denatured (e.g., contacted with guanadine) or unfolded. The unfolded PrP^{Sc} is contacted with a binding partner and the occurrence of binding indicates the presence of PrP^{Sc} in the sample. In another embodiment, PrP^{Sc} and PrP^C are reacted with a labeled antibody that binds both conformations and a conformation that binds only the disease related conformation, and the presence of the disease related conformation is determined by comparing the two.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

ASSAY FOR DISEASE RELATED CONFORMATION OF A PROTEIN

GOVERNMENT RIGHTS

5 The United States Government may have certain rights in this application pursuant to Grant No. AG02132, AG10770, NS22786, NS14069, and NS07219 awarded by the National Institutes of Health.

FIELD OF THE INVENTION

10 This invention relates to the field of bioassays and more particularly to an assay which makes it possible to isolate and detect a disease conformation of a protein present in a native sample also containing a non-disease conformation of the protein.

BACKGROUND OF THE INVENTION

15 Prions are infectious pathogens that cause invariably fatal prion diseases (spongiform encephalopathies) of the central nervous system in humans and animals. Prions differ significantly from bacteria, viruses and viroids. The dominating hypothesis is that no nucleic acid is necessary to allow for the infectivity of a prion protein to proceed.

 A major step in the study of prions and the diseases they cause was the discovery and
20 purification of a protein designated prion protein [Bolton, McKinley et al. (1982) Science 218:1309-1311; Prusiner, Bolton et al. (1982) Biochemistry 21:6942-6950; McKinley, Bolton et al. (1983) Cell 35:57-62]. Complete prion protein-encoding genes have since been cloned, sequenced and expressed in transgenic animals. PrP^C is encoded by a single-copy host gene [Basler, Oesch et al. (1986) Cell 46:417-428] and when PrP^C is expressed it is generally found
25 on the outer surface of neurons. Many lines of evidence indicate that prion diseases results from the transformation of the normal form of prion protein (PrP^C) into the abnormal form (PrP^{Sc}). There is no detectable difference in the amino acid sequence of the two forms. However, PrP^{Sc} when compared with PrP^C has a conformation with higher β -sheet and lower α -helix content [Pan, Baldwin et al. (1993) Proc Natl Acad Sci USA 90:10962-10966; Safar, Roller et al. (1993) J Biol Chem 268:20276-20284]. The presence of the abnormal PrP^{Sc}
30 form in the brains of infected humans or animals is the only disease-specific diagnostic marker of prion diseases.

PrP^{Sc} plays a key role in both transmission and pathogenesis of prion diseases (spongiform encephalopathies) and it is a critical factor in neuronal degeneration [Prusiner (1997) *The Molecular and Genetic Basis of Neurological Disease*, 2nd Edition : 103-143]. The most common prion diseases in animals are scrapie of sheep and goats and bovine spongiform encephalopathy (BSE) of cattle [Wilesmith and Wells (1991) Curr Top Microbiol Immunol 172:21-38]. Four prion diseases of humans have been identified: (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Straussler-Sheinker Disease (GSS), and (4) fatal familial insomnia (FFI) [Gajdusek (1977) Science 197:943-960; Medori, Tritschler et al. (1992) N Engl J Med 326:444-449]. Initially, the presentation of the inherited human prion diseases posed a conundrum which has since been explained by the cellular genetic origin of PrP.

Prions exist in multiple isolates (strains) with distinct biological characteristics when these different strains infect in genetically identical hosts [Prusiner (1997) *The Molecular and Genetic Basis of Neurological Disease*, 2nd Edition:165-186]. The strains differ by incubation time, by topology of accumulation of PrP^{Sc} protein, and in some cases also by distribution and characteristics of brain pathology [DeArmond and Prusiner (1997) *Greenfield's Neuropathology*, 6th Edition:235-280]. Because PrP^{Sc} is the major, and very probably the only component of prions, the existence of prion strains has posed a conundrum as to how biological information can be enciphered in a molecule other than one comprised of nucleic acids. The partial proteolytic treatment of brain homogenates containing some prion isolates has been found to generate peptides with slightly different electrophoretic mobilities [Bessen and Marsh (1992) *J Virol* 66:2096-2101; Bessen and Marsh (1992) *J Gen Virol* 73:329-334; Telling, Parchi et al. (1996) *Science* 274:2079-2082]. These findings suggested different proteolytic cleavage sites due to the different conformation of PrP^{Sc} molecules in different strains of prions. Alternatively, the observed differences could be explained by formation of different complexes with other molecules, forming distinct cleavage sites in PrP^{Sc} in different strains [Marsh and Bessen (1994) *Phil Trans R Soc Lond B* 343:413-414]. Some researchers have proposed that different prion isolates may differ in the glycosylation patterns of prion protein [Collinge, Sidle et al. (1996) *Nature* 383:685-690; Hill, Zeidler et al. (1997) *Lancet* 349:99-100]. However, the reliability of both glycosylation and peptide mapping patterns in diagnostics of multiple prion strains is currently still debated [Collings, Hill et al. (1997) *Nature* 386:564; Somerville, Chong et al. (1997) *Nature* 386:564].

A system for detecting PrP^{Sc} by enhancing immunoreactivity after denaturation is provided in Serban, et al., Neurology, Vol. 40, No. 1, Ja 1990. Sufficiently sensitive and specific direct assay for infectious PrP^{Sc} in biological samples could potentially abolish the need for animal inoculations completely. Unfortunately, such does not appear to be possible with current PrP^{Sc} assays -- it is estimated that the current sensitivity limit of proteinase-K and Western blot-based PrP^{Sc} detection is in a range of 1 µg/ml which corresponds to 10⁴ - 10⁵ prion infectious units. Additionally, the specificity of the traditional proteinase-K-based assays for PrP^{Sc} is in question in light of recent findings of only relative or no proteinase-K resistance of undoubtedly infectious prion preparations [Hsiao, Groth et al. (1994) Proc Natl Acad Sci USA 91:9126-9130] Telling, et al. (1996) Genes & Dev.

Human transthyretin (TTR) is a normal plasma protein composed of four identical, predominantly β-sheet structured units, and serves as a transporter of hormone thyroxine. Abnormal self assembly of TTR into amyloid fibrils causes two forms of human diseases, namely senile systemic amyloidosis (SSA) and familial amyloid polyneuropathy (FAP) [Kelly (1996) Curr Opin Struct Biol 6(1):11-7]. The cause of amyloid formation in FAP are point mutations in the TTR gene; the cause of SSA is unknown. The clinical diagnosis is established histologically by detecting deposits of amyloid *in situ* in biopsy material.

To date, little is known about the mechanism of TTR conversion into amyloid *in vivo*. However, several laboratories have demonstrated that amyloid conversion may be simulated *in vitro* by partial denaturation of normal human TTR [McCutchen, Colon et al. (1993) Biochemistry 32(45):12119-27; McCutchen and Kelly (1993) Biochem Biophys Res Commun 197(2) 415-21]. The mechanism of conformational transition involves monomeric conformational intermediate which polymerizes into linear β-sheet structured amyloid fibrils [Lai, Colon et al. (1996) Biochemistry 35(20):6470-82]. The process can be mitigated by binding with stabilizing molecules such as thyroxine or triiodophenol [Miroy, Lai et al. (1996) Proc Natl Acad Sci USA 93(26):15051-6].

In view of the above points, there is clearly a need for a specific, high flow-through, and cost-effective assay for testing sample materials for the presence of a pathogenic protein including transthyretin and prion protein.

SUMMARY OF THE INVENTION

There are a number of known proteins which exist in two or more conformations. Often, such proteins have a first conformation which is the normal conformation of the protein and a second, disease related conformation. The two proteins are often present
5 together and it is often difficult to determine the presence of the disease related conformation of the protein because (1) it is difficult to obtain binding partners such as antibodies which bind this conformation and (2) the disease related conformation is often present in a relatively low concentration within the sample and relative to the non-disease conformation. The present invention provides assay methods which use various combinations treatments,
10 enzymes and binding partners such as antibodies in order to detect the disease conformation of the protein.

A first assay method of the invention comprises treating a sample suspected of containing a protein which assumes a first conformation and a second, disease related conformation with a compound which hydrolyzes the protein in the first conformation but not
15 the second, disease related conformation of the protein thereby providing a treated sample. The treated sample is then contacted with a binding partner which binds the disease related conformation of the protein. When binding is detected such indicates the presence of the disease related conformation of the protein in the sample. The protein detected may be any protein with two or more conformations and is preferably a PrP protein, a TTR protein or an amyloid protein. When the protein is a PrP protein the compound which hydrolyzes the first
20 conformation of the PrP protein (i.e., PrP^C) is Dispase.

A second embodiment of the assay method of the invention comprises providing a sample suspected of containing a protein which assumes a first confirmation and a second, disease related conformation. That sample is then contacted with a compound which
25 hydrolyzes the protein in the first conformation but not the second, disease related conformation. This results in a treated sample. The treated sample is then subjected to denaturing. More specifically, the unhydrolyzed protein (if any) in the second, disease related conformation is unfolded to some degree in order to allow its epitopes to be more fully exposed. The treated, denatured sample is then contacted with a binding partner which binds
30 to the treated denatured protein, i.e. binds the protein with the exposed epitopes. It is then possible to determine the presence of a protein in the second, disease related conformation by detecting binding of the binding partner.

A third assay method of the invention comprises providing a sample suspected of containing a protein which assumes a first conformation and a second, disease related conformation. The sample is contacted with a first binding partner which binding partner binds only to the first conformation of the protein, i.e. does not bind the second, disease related conformation of the protein. The first binding partner is added in sufficient amounts so as to saturate the sample and bind all and/or substantially all of the available epitopes on any protein in the first conformation present in the sample. Once the first binding partner has been allowed to bind to the available epitopes the sample is then treated with a second binding partner. The second binding partner is different from the first binding partner in that the second binding partner binds to both the first conformation of the protein and the second disease related conformation of the protein. Because all of the available epitopes on the protein of the first conformation are occupied with the first binding partner any detected binding by the second binding partner indicates the presence of the disease related conformation of the protein.

Depending on the steps used in the assay of the invention one of two types of antibodies may be used. Accordingly, both basic types of assays the sample is treated with a compound, e.g. a metalloendopeptidase, which selectively hydrolyzes PrP^C but not PrP^{Sc}. Thereafter, the treated sample can be subjected to two different types of processing, each of which uses a generally different type of antibody.

The first general type of antibody selectively binds to the disease conformation of the protein. For example, antibodies that selectively recognize PrP^{Sc} bind to an epitope on the C-terminus of the protein. When a PrP protein is in its PrP^{Sc} configuration its C-terminus can be bound by antibodies of the type described in U.S. Patent 5,846,533 issued December 8, 1998 - reference is also made to WO 98/37210 which claims to disclose antibodies which bind PrP^{Sc}. Both of these publications are incorporated herein by reference to describe and disclose antibodies and method of making antibodies.

The second general type of antibody binds to both the disease and the non-disease conformations of the protein. For example, antibodies that recognize an epitope on the N-terminus of the PrP protein recognize both PrP^{Sc} and PrP^C following denaturation of the proteins. When the PrP protein is in the PrP^{Sc} configuration the N-terminus is not exposed and as such can not be bound by an antibody. To expose an epitope of the N-terminus the PrP^{Sc} is denatured, e.g. by exposure to guanadine HCl under conditions (pH, temperature,

and time) which causes the PrP^{Sc} to unfold or change its 3-dimensional structure such that a C-terminal epitope is exposed. In this unfolded configuration a wide range of binding partner including commercially available antibodies can be used for detection. Since such antibodies also bind PrP^C all of the PrP^C must be removed, e.g., by selective hydrolysis.

5 An example of an antibody which binds an epitope of the N-terminus is the monoclonal antibody 3F4 produced by the hybridoma cell line ATCC HB9222 deposited on October 8, 1986 in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852 and disclosed and described in U.S. Patent 4,806,627 issued February 21, 1989 - incorporated by reference to disclose antibodies which selectively bind PrP^C in its native
10 form. In addition to antibody other binding partners which bind the non-disease related conformation but not the disease related conformation could be used in the assay of the invention. Antibodies such as 3F4 and others used in the assays described in the examples are commercially available.

 In one embodiment of the invention, one portion of a sample containing two
15 conformations of a protein (e.g. PrP^C and PrP^{Sc}) is reacted with a binding partner (e.g. R1) that binds both conformations, and another portion of the same sample is reacted with a binding partner (e.g. 3F4) that binds only one of the two forms (e.g. PrP^C). The disease related conformation is determined by comparing the two. If the binding partner which binds both conformations shows more binding than the binding partner which binds only one
20 conformation, this shows that both conformations are present in the sample. For example, if R1 binds to more protein than 3F4, PrP^{Sc} is present in the sample. No hydrolysis treatment is needed with this method. However, pretreatment may be used and comparison of the binding may be adjusted for a variety of factors, e.g. binding affinities, comparisons to known samples, hybridization times, variations in signal due to secondary antibodies, etc.

25 An aspect of the invention is to provide an immunoassay which is applicable to assaying samples containing proteins, which samples are suspected of containing a protein which occurs within a native non-disease conformation and a disease related conformation (e.g., PrP protein, β A4 protein and transthyretin).

 Another aspect of the invention is to provide an assay which differentiates between (1)
30 disease related proteins or portions thereof which are not hydrolyzed by limited protease treatment with a protease such as proteinase K (protease resistant proteins, e.g. PrP 27-30)

and (2) disease related proteins which are hydrolyzed by a limited protease treatment with a protease such as proteinase K (e.g., protease-sensitive PrP^{Sc}).

5 An advantage of the present invention is that the immunoassay can quickly and accurately determine the presence of proteins in the disease related conformation (e.g., PrP^{Sc}, β A4 and transthyretin) even though the antibody used in the assay does not bind or has a very low degree of binding affinity for the protein in the disease related conformation and the disease related conformation is present in a lower concentration than the non-disease conformation.

10 A feature of the invention is that the signal obtained can be enhanced by the use of transgenic animals, e.g., mice which are used to detect the presence of a protein in a sample.

Another feature is that time-resolved, dissociation-enhanced fluorescence or a dual wavelength, laser driven fluorometer can be used to enhance sensitivity.

Another advantage is that the assay can detect levels of the disease causing conformation of a protein at a concentration of 1×10^3 particles/ml or less.

15 A specific object is to provide a diagnostic assay for determining the presence of infectious prion protein in variable sample materials obtained or derived from human, primate, monkey, pig, bovine, sheep, goat, deer, elk, cat, dog, mouse, chicken, and turkey tissues and/or body fluids.

20 Another specific object is to provide a diagnostic assay for determining the presence of β A4 protein in variable sample materials obtained or derived from human, primate, monkey, pig, bovine, sheep, goat, deer, elk, cat, dog, mouse, chicken, and turkey tissues and/or body fluids.

25 Another object is to provide a rapid assay for native infectious prion protein in the brains of transgenic and non-transgenic animals injected with sample material potentially containing prions.

Another object is to provide a method to evaluate decontamination procedures by assaying the level of denaturation of pathogenic proteins (e.g., prions or β -sheet β A4) after such treatments.

30 Another advantage is that the process can be carried out without an antibody directly able to recognize an infectious conformation of a protein, and without using a proteinase K step to eliminate the signal of normal (non-disease) isoforms of the protein such as PrP^C.

Another advantage is that in the invented process there is no need for the antibody directly able to recognize a pathogenic conformation of β A4 or transthyretin.

An important feature of the assay is the rapid, cost effective and high flow-through design which can be designed with the capacity to screen 96 samples per day per 96 well
5 plate.

Another aspect of the invention is the diagnostic method to quantitatively detect TTR in the abnormal, amyloid conformation in sample material obtained from human and animal tissues, body fluids, and pharmaceuticals.

The invented process provides a direct, sensitive method to distinguish and quantify
10 the normal and amyloid conformations of TTR in a mixture present in sample materials.

An important object is to provide specific diagnostic assay for pathogenic TTR in variable sample materials obtained or derived from human, primate, monkey, pig, bovine, sheep, deer, elk, cat, dog, and chicken tissues.

Another object is to provide a rapid assay for amyloid form of TTR in transgenic
15 animals.

The specific advantage is that invented assay may detect pathogenic forms of TTR in a mixture with denatured nonpathogenic forms of the same or in a mixture with a soluble form of TTR - for example, detect less than 1×10^3 particles per ml.

These and other objects, advantages, and features of the invented process will become
20 apparent to those skilled in the art upon reading the details of the assay method, antibody development and testing, and transgenic mouse as more fully described below with reference to the attached figures.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

25 Before the present assays and methods are disclosed and described, it is to be understood that this invention is not limited to particular antibodies, proteins, labels, assays or methods as such may, of course vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

30 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein

can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

5 The publications discussed herein are provided solely for the disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of the publications provided are subject to change if it is found that the actual date of publication is different from that provided here.

10

DEFINITIONS

 The term "protein" as used herein is intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term includes naturally occurring proteins and peptides as well as those which are recombinantly or synthetically synthesized.

15 As used in connection with the present invention the term "protein" is specifically intended to cover naturally occurring proteins which occur in at least two different conformations wherein both conformations have the same or substantially the same amino acid sequence but have different three dimensional structures. The two conformations of the protein include at least one conformation which is not related to a disease state and at least one conformation

20 which is related to a disease state -- pathogenic. A specific and preferred example of a protein as used in connection with this disclosure is a PrP protein which includes the non-disease form referred to as the PrP^c form and the disease related form referred as the PrP^{Sc}. Although a prion protein or the PrP^{Sc} form of a PrP protein is infectious and pathogenic, the disease conformation of other proteins is not infectious although it is pathogenic. As used

25 herein, the term pathogenic may mean that the protein actually causes the disease or it may simply mean that the protein is associated with the disease and therefore is present when the disease is present. Thus, a pathogenic protein as used in connection with this disclosure is not necessarily a protein which is the specific causative agent of a disease.

 The terms "pretreatment", "unfolding treatment", and "limited protease treatment" are

30 intended to encompass the descriptions and use of these terms as provided in the respective sections having these headings.

The terms "PrP protein", "PrP" and like are used interchangeably herein and shall mean both the infectious particle form PrP^{Sc} known to cause diseases (spongiform encephalopathies) in humans and animals and the noninfectious form PrP^C which, under appropriate conditions is converted to the infectious PrP^{Sc} form.

5 The terms "prion", "prion protein" and "PrP^{Sc} protein" and the like we used interchangeably herein to refer to the infectious PrP^{Sc} form of PrP, and is a contraction of the words "protein" and "infection." Particles are comprised largely, if not exclusively, of PrP^{Sc} molecules encoded by a PrP gene. Prions are distinct from bacteria, viruses and viroids. Known prions infect animals to cause scrapie, a transmissible, degenerative disease of the nervous system of sheep and goats, as well as bovine spongiform encephalopathy (BSE), or "mad cow disease", and feline spongiform encephalopathy of cats. Four prion diseases known to affect humans are (1) kuru, (2) Creutzfeldt-Jakob Disease (CJD), (3) Gerstmann-Straussler-Scheinker Disease (GSS), and (4) fatal familial insomnia (FFI). As used herein "prion" includes all forms of prions causing all or any of these diseases or others in any animals used - and in particular in humans and domesticated farm animals.

15 The term "PrP gene" is used herein to describe genetic material which expresses proteins including known polymorphisms and pathogenic mutations. The term "PrP gene" refers generally to any gene of any species which encodes any form of a PrP protein. Some commonly known PrP sequences are described in Gabriel et al., Proc. Natl. Acad. Sci. USA 89:9097-9101 (1992), and U.S. Patents 5,565,186; 5,763,740; 5,792,901; and WO97/04814, incorporated herein by reference to disclose and describe such sequences. The PrP gene can be from any animal, including the "host" and "test" animals described herein and any and all polymorphisms and mutations thereof, it being recognized that the terms include other such PrP genes that are yet to be discovered. The protein expressed by such a gene can assume either a PrP^C (non-disease) or PrP^{Sc} (disease) form.

25 The term "binding partner" refers to any molecule which binds the target molecule of interest. Preferably, the binding is of sufficiently high affinity as to make it possible to bind target molecules of interest present in a low concentration, e.g., 1×10^3 particles per ml or less. More preferably the binding partner is selective in binding only the target molecule and not other molecules. Preferred binding partners are antibodies as defined below.

30 The term "antibody" stands for an immunoglobulin protein which is capable of binding an antigen. Antibody as used herein is meant to include the entire antibody as well as any

antibody fragments (e.g. F(ab)', Fab, Fv) capable of binding the epitope, antigen or antigenic fragment of interest. Antibodies for assays of the invention may be immunoreactive or immunospecific for and therefore specifically and selectively bind to a protein of interest e.g., an A4 β amyloid protein or a PrP protein. Antibodies which are immunoreactive and

5 immunospecific for both the native non-disease form and the treated disease form but not for the untreated disease form, (e.g., for both native PrP^C and treated PrP^{Sc} but not native PrP^{Sc}) may be used because the sample is treated to remove i.e., hydrolyze PrP^C. Antibodies for PrP are preferably immunospecific - e.g., not substantially cross-reactive with related materials. Some specific antibodies which can be used in connection with the invention are disclosed in

10 published PCT application WO 97/10505 which is incorporated herein by reference to disclose and describe antibodies. This published PCT application corresponds to USSN 08/713,939. Antibodies disclosed in the PCT application which bind PrP^{Sc} can be used to carry out the basic assay of the present invention when the sample has been treated with dispase sufficiently to hydrolyze all or substantially all of the PrP^C present in the sample.

15 Another useful antibody for binding to PrP^C is the monoclonal antibody 263K 3F4 produced by the hybridoma cell line ATCC HB9222 deposited on October 8, 1986 in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852 and disclosed and described in U.S. Patent 4,806,627 issued February 21, 1989 - incorporated by reference to disclose antibodies which selectively bind PrP^C. The term "antibody" encompasses all types

20 of antibodies, e.g. polyclonal, monoclonal, and those produced by the phage display methodology. Particularly preferred antibodies of the invention are antibodies which have a relatively high degree of affinity for both native PrP^C and treated PrP^{Sc} but a relatively low degree of or substantially no binding affinity for PrP^{Sc}. More specifically, antibodies of the invention preferably have four times or more, more preferably fifteen times or more, and still

25 more preferably 30 times or more binding affinity for both native PrP^C and denatured PrP^{Sc} as compared with the binding affinity for native PrP^{Sc}.

"Purified antibody" refers to that which is sufficiently free of other proteins, carbohydrates, and lipids with which it is naturally associated. Such an antibody "preferentially binds" to a denatured disease conformation of a protein such as the denatured

30 β -sheet conformation of A4 β or PrP^{Sc} protein (or an antigenic fragment thereof), and does not substantially recognize or bind to other antigenically unrelated molecules. A purified antibody of the invention is preferably immunoreactive with and immunospecific for a specific species

and more preferably immunospecific for native PrP^C and for denatured forms of PrP^C and PrP^{Sc} or, alternatively, for native or untreated PrP^{Sc}.

“Antigenic fragment” of a protein (e.g., a PrP protein) is meant a portion of such a protein which is capable of binding an antibody.

5 By “binds specifically” is meant high avidity and/or high affinity binding of an antibody to a specific polypeptide e.g., epitope of a protein, e.g., denatured PrP^{Sc} or denatured A4β protein. Antibody binding to its epitope on this specific polypeptide is preferably stronger than binding of the same antibody to any other epitope, particularly those which may be present in molecules in association with, or in the same sample, as the specific polypeptide of
10 interest e.g., binds more strongly to epitope fragments of a protein such as PrP^{Sc} so that by adjusting binding conditions the antibody binds almost exclusively to an epitope site or fragments of a desired protein such as an epitope fragment exposed by denaturing of PrP^{Sc} and not exposed on native PrP^{Sc}.

By “detectably labeled antibody”, “detectably labeled anti-PrP” or “detectably labeled
15 anti-PrP fragment” is meant an antibody (or antibody fragment which retains binding specificity), having an attached detectable label. The detectable label is normally attached by chemical conjugation, but where the label is a polypeptide, it could alternatively be attached by genetic engineering techniques. Methods for production of detectably labeled proteins are well known in the art. Detectable labels known in the art, but normally are radioisotopes,
20 fluorophores, paramagnetic labels, enzymes (e.g., horseradish peroxidase), or other moieties or compounds which either emit a detectable signal (e.g., radioactivity, fluorescence, color) or emit a detectable signal after exposure of the label to its substrate. Various detectable label/substrate pairs (e.g., horseradish peroxidase/diaminobenzidine, avidin/streptavidin, luciferase/luciferin), methods for labeling antibodies, and methods for using labeled antibodies
25 are well known in the art (see, for example, Harlow and Lane, eds. (Antibodies: A Laboratory Manual (1988) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY)). Europium is a particularly preferred label.

Abbreviations used herein include:

30 CNS for central nervous system;
BSE for bovine spongiform encephalopathy;
CJD for Creutzfeldt-Jacob Disease;

- FFI for fatal familial insomnia;
 GdnHCl for Guanidine hydrochloride;
 GSS for Gerstamnn-Strassler-Scheinker Disease;
 Hu for human;
 5 HuPrP for human prion protein;
 Mo for mouse;
 MoPrP for mouse prion protein;
 SHa for a Syrian hamster;
 SHaPrP for a Syrian hamster prion protein;
 10 Tg for transgenic;
 Tg(SHaPrP) for a transgenic mouse containing the
 PrP gene of a Syrian hamster;
 Tg(HuPrP) for transgenic mice containing the complete human PrP gene;
 Tg(ShePrP) for transgenic mice containing the complete sheep PrP gene;
 15 Tg(BovPrP) for transgenic mice containing the complete cow PrP gene;
 PrP^{Sc} for the scrapie isoform of the prion protein;
 PrP^C for the cellular contained common, normal isoform of the prion protein;
 PrP 27-30 or PrP^{Sc} 27-30 for the treatment or protease resistant form of PrP^{Sc};
 MoPrP^{Sc} for the scrapie isoform of the mouse prion protein;
 20 MHu2M for a chimeric mouse/human PrP gene wherein a region of the mouse PrP
 gene is replaced by a corresponding human sequence which differs from mouse PrP at 9
 codons;
 Tg(MHu2M) mice are transgenic mice of the invention which include the chimeric
 MHu2M gene;
 25 MHu2MP^{Sc} for the scrapie isoform of the chimeric human/mouse PrP gene;
 PrP^{CJD} for the CJD isoform of a PrP protein;
 Prnp^{0/0} for ablation of both alleles of an endogenous prion protein gene, e.g., the
 MoPrP gene;
 Tg(SHaPrP^{+/-})81/Prnp^{0/0} for a particular line (81) of transgenic mice expressing
 30 SHaPrP, +/- indicates heterozygous;

Tg(HuPrP)/Prnp^{0/0} for a hybrid mouse obtained by crossing a mouse with a human prion protein gene (HuPrP) with a mouse with both alleles of the endogenous prion protein gene disrupted;

5 Tg(MHu2M)/Prnp^{0/0} for a hybrid mouse obtained by crossing a mouse with a chimeric prion protein gene (MHu2M) with a mouse with both alleles of the endogenous prion protein gene disrupted;

TTR for transthyretin;

FVB for a standard inbred strain of mice often used in the production of transgenic mice since eggs of FVB mice are relatively large and tolerate microinjection of exogenous
10 DNA relatively well;

[PrP_β] - concentration of prion protein in β-sheet conformation;

[βA4_β] - concentration of βA4 in β-sheet conformation;

[DRC] - concentration of a disease related conformation of a protein.

15 GENERAL ASPECTS OF THE INVENTION

The assay method comprises providing a sample suspected of containing a protein which assumes a first conformation and a second disease related conformation and is capable of detecting a disease conformation of the protein when present in a very low concentration relative to the concentration of other proteins and compounds including the non-disease
20 conformation.

The assay methods disclosed allows one to isolate and detect the presence of a disease related conformation of a protein (e.g., PrP^{Sc}) present in a sample also containing the non-disease related conformation of the protein (e.g., PrP^C). The sample is treated (e.g., contacted with dispase) in a manner which hydrolyzes the PrP^C and not the PrP^{Sc}. The
25 hydrolyzation reaction is stopped (e.g., by the addition of EDTA). The treated sample is contacted with a binding partner (e.g., a labeled antibody which binds PrP^{Sc}) and the occurrence of binding provides an indication that PrP^{Sc} is present. Alternatively the PrP^{Sc} of the treated sample is denatured (e.g., contacted with guanadine) or unfolded so that it has epitopes exposed. The unfolded PrP^{Sc} is contacted with a binding partner (e.g., labeled 3F4)
30 and the occurrence of binding indicates the presence of PrP^{Sc} in the sample.

In accordance with any of the assay embodiments it is preferable to pre-treat the sample being tested to (1) remove as many contaminant proteins as possible; and (2) increase

the concentration of disease related protein in the sample relative to the non-disease related conformation of the protein. For example, the initial sample can be chemically treated with a compound which preferentially degrades or denatures contaminant proteins and/or the relaxed, non-disease form of the protein and/or is exposed to antibodies which preferentially
5 bind to (in order to remove) contaminants and/or non-disease conformation of the protein.

It may be possible to enhance further the sensitivity of various aspects of the invention by concentrating the disease conformation of a protein by adding a compound which selectively binds to the disease conformation to form a complex and centrifuging the sample to precipitate out the complex which is then tested in accordance with the methods described
10 here. Specifics regarding such concentration methods are described in detail in our co-pending application Serial No. 09/026,967 entitled "Process for Concentrating Protein with Disease-Related Conformation".

The different embodiments of the assay of the invention described above are all "direct" types of immunoassays -- meaning that the sample is directly assayed with the labeled
15 antibody either with or without treatment to change the conformation of any disease related conformation proteins present in the sample. An "indirect" assay may also be used. For example, it may be desirable to enhance the number of disease related proteins in the sample (if any) by the use of a transgenic mouse and thereby enhance any signal obtained. To carry out these embodiments of the invention, the sample is first used to inoculate a transgenic
20 mouse which has had its genome modified so that it will develop symptoms of disease when inoculated with proteins in the disease related conformation. After the mice are inoculated, a sufficient period of time is allowed to pass (e.g., 30 days) after which the transgenic animal is sacrificed and a sample such as homogenized brain tissue from the mouse is used in the direct assay described above. The present invention enhances the ability of transgenic mice to
25 detect prions by shortening the period of time which must pass until a determination can be made as to whether the original sample included proteins in the disease related conformation. It would also be possible to use mice of the type disclosed and described in any of U.S. Patents 5,565,186; 5,763,740; or 5,792,901 or to apply epitope tagged PrP as disclosed in U.S. Patent 5,750,361 to affinity purify the PrP^{Sc} from the brain of a Tg mouse and thereafter
30 apply the assay of the present invention. Without the present invention the mouse is inoculated and one must wait until the inoculated mouse actually demonstrates symptoms of the disease. Depending on the mouse, this can take several months or even years. Any of the

assays of the present invention could be used with any transgenic mice such as those described above. The assay could be used well before the mouse developed symptoms of disease thereby shortening the time needed to determine if a sample includes infectious proteins.

5 The assay methodology of the present invention can be applied to any type of sample when the sample is suspected of containing a protein which occurs in at least two conformations. The protein must occur in one conformation which binds to known antibodies, antibodies which can be generated or other specific binding partners. The second conformation must be sufficiently different from the first conformation in terms of its ability
10 to be hydrolyzed by compound (e.g., dispase). In its conceptually simplest form, the invention works best when a compound quickly and complete hydrolyzes the non-disease conformation of the protein without affecting the disease related conformation. However, in reality, a given protein may have more than two conformations. The protein may have more than one non-disease conformation and more than one disease related conformation, (Telling,
15 et al., Science (1996)). The invention is still useful when multiple conformations of non-disease and disease forms of the protein exist -- provided that (1) at least one non-disease conformation differs from at least one disease conformation in terms of its ability to be hydrolyzed by a compound.

As indicated above, the assay of the invention can be used to assay any type of sample
20 for any type of protein, provided the protein includes a non-disease and a disease related conformation. However, the invention was particularly developed to assay samples for the presence of (1) PrP proteins and determine whether the sample included a PrP protein in its disease conformation, i.e., included PrP^{Sc} (2) insoluble forms of β A4 associated with Alzheimer's disease and (3) transthyretin. Accordingly, much of the following disclosure is
25 directed to using the immunoassay of the present invention to detect the presence of either PrP^{Sc} (or to a lesser degree β A4 or transthyretin (TTR)) in a sample -- it being understood that the same general concepts are applicable to detecting disease related conformations of a wide range of different types of proteins.

Europium labeled antibodies used (3F4) have a high binding affinity for PrP^C (non-
30 disease conformation) which comprises an α -helical rich conformation. The antibodies have a low binding affinity for PrP^{Sc} (disease conformation) which comprises a β -sheet rich conformation. The IgG may be obtained from common monoclonal, polyclonal, or

recombinant antibodies, typically recognizing the sequence 90-145 of PrP^C and conformationally unfolded prion protein. Different conformations of recombinant prion protein were chemically crosslinked to polystyrene plates through a glutaraldehyde activation step. The relative affinities of the Eu-labeled IgG with α -helical, β -sheet, and random coil conformation of recombinant Syrian hamster prion protein corresponding to sequence 90-231 were determined by time-resolved, dissociation-enhanced fluorescence in a 96-well polystyrene plate format.

After the labeled antibodies have been provided with sufficient time, temperature and chemical conditions (e.g., pH) to bind to the appropriate proteins present in the respective portions the level of binding of the labeled antibody to protein is determined.

Once a labeled antibody has bound to its target detection may be difficult due to the low concentration of the target molecule in the sample. Different procedures can be used for detection.

Time-resolved, dissociation-enhanced fluorescence and more preferably dual wavelength, laser-driven fluorometers are particularly useful devices - see Hemmilä et al., *Boianalytical Applications of Labeling Technologies* (eds. Hemmilä) 113-119 (Wallas Oy, Turku, Finland, 1995).

These devices make it possible to detect concentrations in an amount in the range of about 1×10^3 particles per ml or less. A high degree of sensitivity is preferred because in most samples the concentration of protein in the disease conformation will be very low. For example, the non-disease conformation of the protein might be present in an amount of about 1×10^8 particles/ml while the disease conformation of the protein is only present in an amount of 1×10^4 particles/ml.

The assay can be used to test for the presence of the disease conformation of a given protein within any type of sample. Some of the most typical samples to be tested include pharmaceuticals which include components which are derived from living mammals or use materials derived from living mammals in their processing. It would also be desirable to test organs for transplantation and food items such as beef which was suspected of containing infectious prions. The invention could be used for testing for the presence of the disease conformation of one or more types of proteins such as infectious PrP^{Sc} in pharmaceuticals, cosmetics, biopsy or autopsy tissue, brain, spinal cord, peripheral nerve, muscle, cerebrospinal fluid, blood and blood components, lymph nodes, and in animal or human-derived cultures

infected or potentially infected by disease forms of proteins such as prions. The brains of cows suspected of being infected with prions (i.e., BoPrP^{Sc}) could be tested to determine if the cows can be safely used for human consumption.

5

TREATMENT - GENERAL

An assay of the invention can use all or any of three basic types of treatment which are defined above. The treatments are (1) pretreatment, (2) unfolding treatment and (3) hydrolysis treatment. In general the conditions for pretreatment are gentle, those for unfolding treatment moderate and those for hydrolysis treatment are harsh. Each type of treatment can employ the same means (e.g. proteases, time, pH, temperature, etc.) but employs each to a different degree, e.g. higher concentration, longer time, higher temperature. However, the hydrolysis treatment must employ a compound which selectively hydrolyzes only the non-disease conformation and not the disease conformation.

15

PRETREATMENT

Before carrying out treatment or antibody testing of the sample it may be desirable to subject the sample to pretreatment. The pretreatment is carried out in order to destroy or remove unrelated proteins as well as some of the non-disease form of the protein present within the sample. Examples of pretreatment methodology include producing a column which includes antibodies bound to support surfaces which antibodies bind to the non-disease conformation of the protein thereby removing as much of the non-disease conformation of the proteins possible. Antibodies which bind unrelated but common proteins can also be used. Alternatively, the sample can be subjected to physical treatment such as long term hydrostatic pressure or temperature alone or in combination with chemicals such as acids or alkalines as indicated above to destroy proteins present in the sample which proteins are not related to those being assayed for or are in the non-disease conformation. In some instances proteins in the non-disease and disease conformation will be destroyed. However, a higher relative percentage of the proteins in the non-disease conformation will be destroyed because these proteins are initially in a looser conformation which is more vulnerable to destruction. Thus, the pretreatment methodology results in a sample which includes a relatively lower concentration of the non-disease conformation of the protein relative to the concentration of the disease conformation of the protein. Further, the pretreated sample will have a lower

concentration of unrelated proteins. This increases the sensitivity of the assay making it possible to detect lower concentrations of the disease conformation of the protein. Removal of proteins is preferred over destruction of such in that destruction will decrease sensitivity if the disease conformation is destroyed. A particularly useful pretreatment method is disclosed in our patent application Serial No. 09/026,967 entitled "Process for Concentrating Protein with Disease-Related Conformation".

UNFOLDING TREATMENT

The unfolding treatment denatures the protein but does not hydrolyze proteins of interest and can include exposing the proteins to any physical and/or chemical means which causes the protein which is originally present in a tightened, disease related conformation (e.g., PrP^{Sc}) to assume a more relaxed conformation which has a higher degree of binding affinity for any binding partner such as antibodies (e.g., expose an N-terminal epitope of PrP^{Sc}). In general, the unfolding treatment involves subjecting the protein to some means which causes epitopes on the protein which were not previously exposed or partially exposed to become exposed or become more exposed so that an antibody or other binding partner can more readily bind to the newly exposed epitope.

Methods used for unfolding treatment may include: (1) physical, such as hydrostatic pressure or temperature, (2) chemical, such as acidic or alkaline pH, chaotropic salts, denaturing detergents, guanidine hydrochloride and proteinases such as Proteinase K and (3) combinations of above. The concentration, temperature and time must be considered so as to obtain the desired effect, e.g. unfolding but not hydrolysis.

The treatment time will vary depending on the treatment used but should be carried out for sufficient time to obtain the desired effect, e.g. for unfolding treatment to expose new binding sites but not so long as to completely denature or hydrolyze the protein. When carrying out unfolding treatment on PrP proteins without chemical treatment the temperature is raised to about 40°C to about 80°C for a time sufficient to obtain the desired amount of unfolding of PrP^{Sc}. The temperature can be lower and the time shorter if the pH is raised to 12 or 13.

30

HYDROLYSIS TREATMENT

The hydrolysis treatment is a lytic treatment which is the most important treatment method used in one embodiment of the assays of the invention. After a sample has been subjected to the pretreatment treatment it is subjected to the hydrolysis treatment. This treatment will destroy or hydrolyze all or substantially all protein in the sample which is in the non-disease conformation and not hydrolyze the protein in the disease conformation. The hydrolysis treatment is preferably via an enzyme such as a hydrolase that acts on peptide bonds, preferably a neutral protease, more preferably a metalloendopeptidase, and most preferably dispase or leucostoma peptidase A. The proteases used in the method of the invention may be used alone, in combination, or in conjunction with enzymes having similar but distinct activity such as a carbohydrase, e.g. collagenase, amylase, or alkaline serine protease. The concentration of the treating compounds as well as the time and temperature will vary with the protein being treated and end result to be obtained. For example, with PrP the treatment is carried out to hydrolyze all or substantially all non-PrP^C present, but not hydrolyze PrP^{Sc} present. The object of this treatment is to hydrolyze as much non-disease protein as possible (preferably all) while hydrolyzing as little (preferably none) disease related protein as possible. The treatment is preferably designed such that it can be quickly and completely stopped at any given time. For example, hydrolysis of PrP^C with dispase or other related proteases can be stopped by adding EDTA.

The following list of enzymes are preferred compounds of the method of the invention:

Enzyme	Biological Source
Dispase	<i>Bacillus polymyxa</i>
Atrolysin A, B, C, E and F	Western diamondback rattlesnake <i>Crotalus atrox</i>
Envelysin	Various member of the class <i>Echinoidea</i>
Thimet oligopeptidase	Related to Saccharolysin from <i>Saccharomyces cerevisiae</i> .
Matrilysin	Rat uterus
Vibriolysin	<i>Vibrio proteolyticus</i> (formerly <i>Aeromonas proteolytica</i>)
Coccolysin	<i>Streptococcus thermophilus</i>
Mycolysin	<i>Streptomyces griseus</i>
Meprin A	rat and mus kidney and intestinal brush border
Astacin	The crayfish <i>Astacus fluviatilis</i>
Aureolysin	<i>Staphylococcus aureus</i>
Leishmanolysin	Various species of <i>Leishmania</i> protozoans
Peptidyl-Asp Metalloendopeptidase	<i>Psuedomonas fragi</i>
Autolysin	<i>Chlamydomonas reinhardtii</i>

	Deuterolysin	<i>Penicillium roqueforti</i> ; species variants include <i>Penicillium caseicolum</i> , <i>Aspergillus sojae</i> , and <i>Aspergillus oryzae</i> .
	Bothrolysin	Venom of jararaca snake <i>Bothrops jararaca</i>
	Stromelysin 1 and 2	Human rheumatoid synovial fibroblasts
	Bacillolysin	<i>Bacillus subtilis</i> ; species variants include <i>Bacillus amyloliquefaciens</i> , <i>Bacillus megaterium</i> , <i>Bacillus mesentericus</i> , <i>Bacillus cereus</i> , and <i>Bacillus stearothermophilus</i>
5	Thermolysin	<i>Bacillus thermoproteolyticus</i> ; species variants include <i>Micrococcus caseolyticus</i> and <i>Aspergillus oryzae</i> .
	Acromonolysin	Honey fungus <i>Armillaria mellea</i>
	Leucolysin	Venom of the western cottonmouth moccasin snake, <i>Agkistrodon piscivorus</i>
	Mycolysin	<i>Streptomyces griseus</i> , <i>Streptomyces naraensis</i> , and <i>Streptomyces cacaoi</i>
	Pseudolysin	<i>Pseudomonas aeruginosa</i>
10	Peptidyl-Lys Metalloendopeptidase	<i>Pseudomonas fragi</i>
	Aureolysin	<i>Staphylococcus aureus</i>
	Neprilysin	Widely distributed in mammal tissues, including brain, liver and lung; abundant in kidney brush border membrane
	β -lytic Metalloendopeptidase	<i>Achromobacter lyticus</i> and <i>Lysobacter enzymogenes</i>
	Peptidyl-Asp Metalloendopeptidase	<i>Pseudomonas fragi</i>
15	Ophiolysin	Venom of the King Cobra <i>Ophiophagus hannah</i> .
	Pitrilysin	<i>Escheria coli</i>
	Insulysin	mammals and <i>Drosophila melanogasters</i>
	Serralysin	<i>Pseudomonas aeruginosa</i> ; species variants include <i>Escheria freundii</i> , <i>Serratia marcescens</i> , and <i>Erwinia chrysanthemi</i> .

- 20 The method of the invention is not limited to these enzymes, and thus other enzymes predicted by those skilled in the art to function in the method of the invention may be used.

BINDING PROTEINS TO SUPPORT SURFACES

- 25 The method of chemical or affinity coupling of PrP protein to the plastic support are generally described in available literature and may vary. The antibodies used in the diagnostic assay are polyclonal, monoclonal or recombinant Fab and need to be species specific with preferential binding to the native PrP^C or denatured form of PrP^{Sc} with preferably at least 4-fold lower reactivity with infectious PrP^{Sc}, assuming the same amount of the antigen.

USING THE ASSAY TO DETECT PRIONS (PrP^{Sc})

One aspect of the invention is a two step process to diagnose prion disease by quantitatively measuring the native infectious form of PrP^{Sc} protein in sample material or in the brains of susceptible animals inoculated with such material. The sample is preferably pretreated to remove as much unrelated and non-disease protein as possible. The pretreated sample is first subjected to hydrolysis treatment and then crosslinked to the plastic support.

To measure the concentration of PrP^{Sc} when it is much less than PrP^C, the detection system has to have extreme sensitivity and a linear range of at least 10⁴. The assay described herein can readily detect PrP^{Sc} at a concentration of (approximately) 50 pg/ml using Europium-labeled IgG. Assuming 10⁵ - 10⁶ PrP^{Sc} molecules per ID₅₀ unit the present assay can readily detect 5 x 10² - 5 x 10³ ID₅₀ units per ml.

The assay can detect PrP^{Sc} in mixtures (by direct method) where the concentration of PrP^{Sc} is less than 1% of the concentration of PrP^C. Additional sensitivity can be achieved by immunoprecipitation, using a sandwich format for a solid state assay, differential centrifugation with detergent extraction to remove PrP^C, the indirect transgenic animal method or combinations of these methods. A conservative estimate is that such procedures should allow measurement of between 5 and 50 ID₅₀ units per ml or less conservatively to measure between 0.1 and 0.01 ID₅₀ units per ml. Such measurements would provide a rapid, "positive" means of establishing biological sterility which is the "absence" of infectivity.

20

ANTIBODIES

Method of generating antibodies are generally known to those skilled in the art. In that the disease form is often in a tighter configuration than the non-disease form, with less epitopes exposed, one can readily generate antibodies which bind only to the non-disease form of the protein or the treated disease form. For example, antibodies detecting treated forms of PrP^{Sc} protein and PrP^C protein may be generated by immunizing rabbits or mice with α -helical conformations of recombinant PrP, native PrP^C from animal brains, synthetic peptides in α -helical or random coil conformations, or against denatured PrP^{Sc} or PrP 27-30. Only antibodies with affinity at least 4 fold higher for PrP^C (or denatured conformation of PrP^{Sc} of the same species) as compared to their affinity for PrP^{Sc} should be selected. The method of antibody generation, purification, labeling and detection may vary.

30

The IgG or Fab's may be purified from different sources by affinity HPLC using protein A column and Size exclusion HPLC. The purified antibodies may be labeled with Europium and detected by time resolved fluorescence. The antibody binding to different conformations of PrP protein may be measured by time-resolved, dissociation-enhanced fluorescence. However, the system of detection of PrP-bound IgG on solid support in situ or in solution may vary. Further, it is possible to use direct or indirect immunological methods including direct radiolabels, fluorescence, luminescence, avidin-biotin amplification, or enzyme-linked assays with color or luminescent substrates.

An antibody which can be used in the invention is disclosed in US 4,806,627, issued February 21, 1989, disclosing monoclonal antibody 263K 3F4, produced by cell line ATCC HB9222 deposited on October 8, 1986, which is incorporated herein by reference. The cell line producing the antibody can be obtained from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852.

In general, scrapie infection fails to produce an immune response, with host organisms being tolerant to PrP^{Sc} from the same species. Antibodies which bind to either PrP^C or PrP^{Sc} are disclosed in U.S. Patent 5,846,533 issued Dec. 8, 1998. Any antibody binding to PrP^C and not to PrP^{Sc} can be used, and those skilled in the art can generate such using known procedures, e.g., see methods of producing page display antibody libraries in US 5,223,409. Polyclonal anti-PrP antibodies have though been raised in rabbits following immunization with large amounts of formic acid or SDS-denatured SHaPrP 27-30 [Bendheim, Barry et al. (1984) Nature 310:418-421; Bode, Pocchiari et al. (1985) J Gen Virol 66:2471-2478; Safar, Ceroni et al. (1990) Neurology 40:513-517]. Similarly, a handful of anti-PrP monoclonal antibodies against PrP 27-30 have been produced in mice [Barry and Prusiner (1986) J Infect Dis 154:518-521; Kascsak, Rubenstein et al. (1987) J Virol 61:3688-3693]. These antibodies were generated against formic acid- or SDS-denatured PrP 27-30 and are able to recognize native PrP^C and treated or denatured PrP^{Sc} from both SHa and humans equally well, but do not bind to native PrP^{Sc}. Not surprisingly, the epitopes of these antibodies were mapped to regions of the sequence containing amino acid differences between SHa- and MoPrP [Rogers, Yehiely et al. (1993) Proc Natl Acad Sci USA 90:3182-3186].

It is not entirely clear why many antibodies of the type described in the above cited publications will bind to PrP^C and treated or denatured PrP^{Sc} but not to native PrP^{Sc}. Without being bound to any particular theory it is suggested that such may take place because epitopes

which are exposed when the protein is in the PrP^C conformation are unexposed or partially hidden in the PrP^{Sc} configuration - where the protein is relatively insoluble and more compactly folded together.

For purposes of the invention an indication that no binding occurs means that the equilibrium or affinity constant K_a is 10^6 l/mole or less. Further, binding will be recognized as existing when the K_a is at 10^7 l/mole or greater, preferably 10^8 l/mole or greater. The binding affinity of 10^7 l/mole or more may be due to (1) a single monoclonal antibody (i.e., large numbers of one kind of antibodies) or (2) a plurality of different monoclonal antibodies (e.g., large numbers of each of five different monoclonal antibodies) or (3) large numbers of polyclonal antibodies. It is also possible to use combinations of (1) - (3). Selected preferred antibodies will bind at least 4-fold more avidly to the treated or denatured PrP^{Sc} forms of the protein when compared with their binding to the native conformation of PrP^{Sc}. The four fold differential in binding affinity may be accomplished by using several different antibodies as per (1) - (3) above and as such some of the antibodies in a mixture could have less than a four fold difference.

A variety of different types of assays of the invention may be used with one or more different antibodies. Those skill in the art will recognize that antibodies may be labeled with known labels and used with currently available robotics, sandwich assays, electronic detectors, flow cytometry, and the like.

DISEASES ASSOCIATED WITH INSOLUBLE PROTEINS

Much of the disclosure and the specific examples provided herein relate to the use of the assay in connection with determining the presence of PrP^{Sc} in the sample. However, as indicated above, the assay of the invention can be applied to determining the presence of any protein which assumes two different conformational shapes, one of which is associated with the disease. The following is a non-limiting list of diseases with associated insoluble proteins which assume two or more different conformations.

	<u>Disease</u>	<u>Insoluble Proteins</u>
	Alzheimer's Disease	APP, A β peptide, α 1-antichymotrypsin, tan, non-A β component
5	Prion diseases, Creutzfeld Jakob disease, scrapie and bovine spongeform encephalopathy	PrP ^{Sc}
	ALS	SOD and neurofilament
	Pick's disease	Pick body
	Parkinson's disease	Lewy body
10	Diabetes Type 1	Amylin
	Multiple myeloma--plasma cell dyscrasias	IgGL-chain
	Familial amyloidotic polyneuropathy	Transthyretin
	Medullary carcinoma of thyroid	Procalcitonin
	Chronic renal failure	β_2 --microglobulin
15	Congestive heart failure	Atrial natriuretic factor
	Senile cardiac and systemic amyloidosis	Transthyretin
	Chronic inflammation	Serum amyloid A
	Atherosclerosis	ApoA1
	Familial amyloidosis	Gelsolin

20

It should be noted that the insoluble proteins listed above each include a number of variants or mutations which result in different strains which are all encompassed by the present invention. Known pathogenic mutations and polymorphisms in the PrP gene related to prion diseases are given below and the sequences of human, sheep and bovine are given in

25 US 5,565,186, issued October 15, 1996.

MUTATION TABLE

	Pathogenic human mutations	Human Polymorphisms	Sheep Polymorphisms	Bovine Polymorphisms
	2 octarepeat insert	Codon 129 Met/Val	Codon 171 Arg/Glu	5 or 6 octarepeats
5	4 octarepeat insert	Codon 219 Glu/Lys	Codon 136 Ala/Val	
	5 octarepeat insert			
	6 octarepeat insert			
	7 octarepeat insert			
	8 octarepeat insert			
10	9 octarepeat insert			
	Codon 102 Pro-Leu			
	Codon 105 Pro-Leu			
	Codon 117 Ala-Val			
	Codon 145 Stop			
15	Codon 178 Asp-Asn			
	Codon 180 Val-Ile			
	Codon 198 Phe-Ser			
	Codon 200 Glu-Lys			
	Codon 210 Val-Ile			
20	Codon 217 Asn-Arg			
	Codon 232 Met-Ala			

It should also be noted that such proteins have two different 3-dimensional conformations with the same amino acid sequence. One conformation is associated with disease characteristics and is generally insoluble whereas the other conformation is not associated with disease characteristics and is soluble. The methodology of the present invention is not limited to the diseases, proteins and strains listed.

DETECTING THE β -SHEET FORM OF β A4

One aspect of the invention involves a two step process to diagnose Alzheimer's disease based on the presence of a constricted form of a protein (β A4 amyloidosis) by quantitatively measuring β -sheet form of β A4 protein in sample material, e.g., in the brain or body fluids. The sample is divided into two aliquots. The first aliquot is crosslinked to a solid plastic (long chain polymeric material) support in native conformation through a chemical activation step under the nondenaturing conditions. The second portion of the sample is first subjected to unfolding treatment and then crosslinked to the plastic support. Both portions of the sample material react in situ with the labeled antibodies that

preferentially recognize soluble β A4 or unfolding treatment β A4 of the human or a given animal species. The amount of the antibody bound to unfolded or native conformations of β A4 protein is recorded by the signal of the labeled secondary antibody. The excess of the signal obtained with the unfolding treated sample compared to that expected change in the signal obtained with the native α -helical conformation of β A4 protein is the measure of the amount of β -sheet structured β A4 in the original sample. The formula developed for calculation of β A4 content is provided above in connection with the calculation of PrP^{Sc} content.

The diagnosis of β A4 amyloidosis (Alzheimer's disease) is established by three procedures: (1) measurement of denatured sample alone and by detecting the increase in the total β A4 amount (concentration) in the examined sample above the background levels of soluble β A4 obtained from normal controls; (2) calculation of the ratio between unfolding treated versus native signal for a given antibodies (protein index) - for example values higher than 2 for monoclonal antibody 6F3D and europium labeled secondary antibody; (3) evaluation of the change of the denatured sample signal over that expected change in the signal for α -helical conformation of β A4 as a measure of the amount of infectious β -sheet structured β A4 in the original sample. The formula developed for calculation of β A4 content is provided above. The particular strain of β A4 can also be determined using the same methodology described above to determine the strain of PrP^{Sc} in a sample.

The invention provides a direct diagnostic method for detecting the presence pathogenic forms of β A4 protein in pharmaceuticals, biopsy or autopsy tissue, brain, spinal cord, peripheral nerves, muscle, cerebrospinal fluid, blood and blood components, lymph nodes, and in animal- or human-derived cultures expressing or potentially expressing β A4 protein. The invention also makes it possible to follow the α -helix-to- β -sheet conformational transition of β A4 protein, or its fragments of synthetic or recombinant origin, and to provide a method to screen compounds for their ability to stabilize the normal soluble conformation of β A4 protein and thus prevent conversion into pathogenic insoluble and β -sheet-structured β A4 protein.

Typical methods of sample denaturation include: (1) physical, such as hydrostatic pressure or temperature, (2) chemical, such as acidic or alkaline pH, chaotropic salts, or denaturing detergents, and (3) combination of above. Methods of chemical or affinity coupling of β A4 protein to a plastic support are described in available literature and may vary.

Antibodies used in the diagnostic assay may be polyclonal, monoclonal or recombinant Fab and must be species specific with preferential binding to the soluble or denatured form of β A4 with preferably at least a 2-fold difference in reactivity between α -helical and β -sheet structured β A4, assuming the same amount of antigen.

5 Methods of sample attachment to the plastic support may vary and may be covalent or non-covalent as described in available literature. The sensitivity of the assay described in the examples may be increased by using high-affinity antibodies, sandwich formate, immunoprecipitation, or differential centrifugation. However, only the antibodies with an affinity at least a 2 fold for unfolding treated as compared to the native β -sheet conformation
10 of β A4 of the same species shall be used for the diagnostic assay. Methods of antibody generation, purification, labeling and detection may vary. The antibody binding to different conformations of β A4 protein was measured by time-resolved, dissociation-enhanced fluorescence. However, the system of detection of β A4-bound IgG on solid support in situ or in solution may vary and may use direct or indirect immunological methods including direct
15 radiolabels, fluorescence, luminescence, avidin-biotin amplification, or enzyme-linked assays with color or luminescent substrates.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art
20 with a complete disclosure and description of how to make and use assays of the present invention, and are not intended to limit the scope of what the inventors regard as their invention, nor are they intended to represent or imply that the experiments below are all of or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations
25 should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees centigrade, and pressure is at or near atmospheric.

EXAMPLE 1

30 DETECTION OF PrP^{Sc} IN HAMSTER BRAIN

To determine the levels of PrP^{Sc} in affected hamsters, a prion infected hamster and a normal hamster were each sacrificed and their brains removed. A 10% (w/v) homogenate of

each of the brains was prepared by dispersing the brain tissue in PBS. The brain homogenate was then subjected to a low speed centrifugation of 500xg for 15 minutes to separate the suspended proteins from unwanted cellular debris. The total protein concentration of the supernatant (S1) was measured using a BCA Protein Assay (Pierce) and the concentration of
5 each brain homogenate was adjusted with PBS to 3.5 mg/ml. A portion of the homogenate was saved to serve as a control of total brain proteins.

The metalloendopeptidase dispase (Worthington) was added to the remainder of each sample at an enzyme to protein ratio of 1:35. The homogenates were digested with 100 μ g/ml dispase for 60 min at 37°C in the presence of either 0.15% Zwittergent 3-12
10 (Calbiochem), or 0.2% or 2% of sodium dodecyl sarcosinate (Sarkosyl). A sample without the dispase was also placed at 37°C to serve as a digestion control. Following digestion the dispase was inactivated by the addition of 50 mM EDTA.

As a control, Proteinase K digestion was done on each untreated S1 sample with 20 μ g/ml at a ratio of enzyme to protein of 1:50 (Bolton, 1982) in the presence of either 0.15%
15 Zwittergent, 0.2% or 2% Sarkosyl. These reactions were terminated by the addition of 2 mM PMSF.

The digested samples were centrifuged at 100,000 x g for 1 hour. The pellet containing all the insoluble proteins was resuspended in a minimal volume of PBS and 0.15% Zwittergent. Subsequently, both the digested samples and the whole brain homogenate
20 samples were sonicated in a water bath sonicator for 20 minutes to denature the protein remaining after digestion.

Each sample was then analyzed for PrP protein content by immunoblot. A 100 μ l aliquot of each sample was placed in a 1.5 ml eppendorf tube with an equal volume of sample loading buffer (1X=50 mM TrisCl, pH 6.8; 100 mM DTT; 2% SDS; 0.1 % bromophenol blue;
25 and 10% glycerol). In addition, an aliquot of b-mercaptoethanol can be added to ensure denaturation of the proteins on the gel. The samples were run on a 10% polyacrylamide gel, molar ratio of bisacrylamide:acrylamide of 1:29, for 15 V/cm for about 4 hours. The samples were each boiled for 10 minutes, and 15 μ l of each sample was loaded onto the gel. For a more detailed description of protein separation via PAGE, see Schagger and Von Jagow,
30 *Nature* 166:368-379 (1987) and Laemmli, U.K. (1970) *Nature* 227, 680-685, which are both incorporated herein in their entirety.

The gel was removed from the PAGE apparatus, and transferred onto uncharged nylon (Amersham) using an electroblotting apparatus. The gel and the nylon were sandwiched between pieces of Whatman 3MM paper soaked in a transfer buffer containing Tris, glycine, SDS and methanol. The sandwich was placed between graphite plate electrodes, with the nylon on the anodic side. A current of 0.65 mA/sq. cm was applied for 1.5-2 hours. Following transfer, the gel was stained with coomassie blue to be sure the transfer was complete.

The nylon filter was placed in a heat-sealable plastic bag, and 0.1 ml of blocking solution were added per square cm of filter. The blocking solution contained 5% (w/v) nonfat dry milk (Carnation); 0.01% antifoam A; and 0.02% sodium azide in PBS. After 1 hour shaking at room temperature, the monoclonal antibody 3F4 was added in a 1:100 dilution, and the filter incubated for 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody were removed, and the filter washed three times.

The filter was incubated with an anti-Ig secondary antibody in blocking solution for 1-2 hours at room temperature. The secondary antibody was radiolabeled to allow immunoblot detection. For the radiolabeled I¹²⁵ probe, approximately 10⁴ cpm of the reagent was added per square centimeter of filter. After incubation, the filter was washed several times in PBS, each wash being about 10 minutes in length. The filter was placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C.

The results of the immunoblot were as follows:

lane 1 normal hamster S1

Result: strong band at about 33-35 Kd

lane 2 normal hamster S1 at 37°C

Result: strong band at about 33-35 Kd

lane 3 normal hamster S1 digested with 100µg/ml dispase, 0.15% Zwittergent

Result: no band

lane 4 normal hamster S1 digested with 100µg/ml dispase, 0.2% Sarkosyl

Result: no band

lane 5 normal hamster S1 digested with 100µg/ml dispase, 2% Sarkosyl

Result: no band

- lane 6 normal hamster S1 digested with 20 μ g/ml Proteinase K, 0.2% Sarkosyl
Result: no band
- lane 7 normal hamster S1 digested with 20 μ g/ml Proteinase K, 2% Sarkosyl
Result: no band
- 5 lane 8 prion-infected hamster S1
Result: strong band at about 33-35 Kd
- lane 9 prion-infected hamster Si at 37°C
Result: strong band at about 33-35 Kd
- 10 lane 10 prion-infected hamster S1 digested with 100 μ g/ml dispase, 0.15% Zwittergent
Result: very strong band at about 33-35 Kd
- lane 11 prion-infected hamster S1 digested with 100 μ g/ml dispase, 0.2% Sarkosyl
Result: very strong band at about 33-35 Kd
- lane 12 prion-infected hamster S1 digested with 100 μ g/ml dispase, 2% Sarkosyl
Result: very strong band at about 33-35 Kd
- 15 lane 13 prion-infected hamster S1 digested with 20 μ g/ml protein kinase, 0.2% Sarkosyl
Result: very strong band at about 27-30 Kd
- lane 14 prion-infected hamster S1 digested with 20 μ g/ml protein kinase, 2% Sarkosyl
Result: very strong band at about 27-30 Kd

20

These results showed that normal hamster brain PrP^C was not detected after digestion with Dispase or Proteinase K, while samples from the prion infected brain showed a very strong signal of protease resistant protein. The prion infected samples digested with proteinase K showed an expected shift in the size of the molecular weight corresponding to the digestion of

25 the n-terminus of the protein. The digestion with Dispase showed no shifting in the molecular weight. This finding showed that the Dispase is selective for the normal conformation of the protein.

EXAMPLE 2

30

DETECTION OF PrP^{Sc} IN MOUSE BRAIN

To determine the levels of PrP^{Sc} in affected mice, a prion infected mouse and a normal mouse are each sacrificed and their brains removed. A 10% (w/v) brain homogenate from

normal and prion infected mice is prepared in PBS. After a low speed centrifugation at 500 X g for 15 min, the total protein in the supernatant (S1) is measured using spectrophotometric assays, and the concentration is adjusted to 2.5 mg/ml with PBS.

The samples are digested with 500 U/ml Leucolysin for 45 min at 37°C in the presence of 2% Sarkosyl. The digestion is stopped by the addition of 50 mM EDTA. An aliquot of the proteins obtained in S1 both before and after the leucolysin digestion are electrophoresed at 4°C on an 8% polyacrylamide slab gel as described in the Laemmli reference but in the absence of SDS and 2-mercaptoethanol. This allows the nondenatured proteins to migrate through the polyacrylamide while preserving the native structure of the protein. Once immobilized in the polyacrylamide, the proteins of the gel are then transferred to nitrocellulose for protein detection. Transfer may occur as in Example 1, or a semi-dry transfer apparatus may be used (Reference Maniatis).

The digested and undigested sample of both infected and normal mouse are detected in Western blot as described in Example 1, but using a monoclonal antibody that recognizes the native PrP^{Sc} form of the protein. Such antibodies are described in WO 98/37411 published August 27, 1998 and U.S. Patent 5,846,533 issued December 8, 1998, each of which is incorporated herein by reference in their entirety. Preferably, the antibody used is the PrP^{Sc}-specific antibody R1. The antibodies are added at a concentration of about 1:100 to 1:200, depending on the antibody used and the amount of protein predicted to be immobilized on the nitrocellulose.

The nitrocellulose is placed in a heat-sealable plastic bag, and 0.1 ml of blocking solution is added per square cm of filter. The blocking solution contains 5% (w/v) nonfat dry milk (Carnation); 0.01% antifoam A; and 0.02% sodium azide in PBS, and Tween 20 added to a final concentration of 0.02%. The Tween 20 is a gentle detergent that further aids in reducing the background. After 1 hour shaking at room temperature, the R1 antibody is added in a 1:100 dilution, and the filter is incubated for 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody are removed, and the filter is washed three times.

The filter is incubated with an anti-Ig secondary antibody in blocking solution for 1-2 hours at room temperature. The secondary antibody is enzyme-conjugated with horseradish peroxidase to allow immunoblot detection. The secondary antibody is added at a much more dilute level than the primary antibody, from 1:500 to 1:2000 dilution. After incubation, the

filter is washed several times in PBS, each wash being about 10 minutes in length. The filter is then placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C .

The resulting blot will have a band corresponding to PrP^{Sc} in the sample of the treated, prion infected mouse brain, both before and after digestion with Leucolysin. The sample of the normal mouse brain protein will show a low level band due to binding of the PrP^{Sc} . The lane with the normal brain sample subjected to hydrolysis with Leucolysin, however, will not have a detectable band, since the Leucolysin will have hydrolyzed the PrP protein in the sample.

EXAMPLE 3

DETECTION OF PrP^{Sc} IN COW BRAIN

A 10% (w/v) brain homogenate from normal and prion infected cows is resuspended in 1 L of 25 mM Tris-HCl, pH 8.0, 5 mM EDTA (buffer A). This is centrifuged at $10,000 \times g$ for 20 min, and the supernatant containing soluble cytoplasmic proteins is discarded. The pellet is resuspended in 1 ml of buffer A, passed through a cell disrupter twice (Microfluidics International, model MF110), and centrifuged at $30,000 \times g$ for 1 h, after which the supernatant is discarded and the pellet is washed once in buffer A and centrifuged again at $30,000 \times g$ for 1 hour. At this stage the pellet could be stored at -20°C prior to hydrolysis.

The β -Lytic Metalloendopeptidase digestion is done at an enzyme to protein ratio of 1:40. The protein is digested with $100 \mu\text{g/ml}$ β -Lytic Metalloendopeptidase (Sigma) for 75 min at 40°C in a buffered pH 8.0 solution containing 0.2% Sarkosyl. The digestion is stopped by the addition of 50 mM EDTA. Following the hydrolysis of the PrP^{C} conformation of the prion protein, the PrP^{Sc} can be denatured to allow the 3F4 antibody to recognize its epitope. Both the normal and prion infected samples are treated with a denaturation solution of 6M guanidine HCl. The denaturation solution is prepared by diluting 10X buffer (250mM HEPES (pH 7.9); 30 mM MgCl_2 ; 40 mM KCl) with five volumes of distilled water. An appropriate amount of the guanidine HCl is added, and the solution is brought to 1X using distilled water. Finally, dithiothreitol is added to a final concentration of 1mM. The hydrolyzed samples are subjected to treatment with the denaturation solution for 30 minutes at 4°C .

The samples are then loaded onto 10% polyacrylamide gels as described in Example 1, and transferred to nylon membrane. The nylon membrane is placed in a heat-sealable

plastic bag, and 0.1 ml of blocking solution is added per square cm of filter. The blocking solution contains 5% (w/v) nonfat dry milk (Carnation) and 0.02% sodium azide in PBS. After 1 hour shaking at room temperature, the monoclonal antibody 3F4 is added in a 1:200 dilution, and the filter incubated for 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody are removed, and the filter washed three times. The filter is incubated with an anti-Ig secondary antibody in blocking solution for 1-2 hours at room temperature, the filter washed several times in PBS, and placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C.

The determination of prion infection is based on a comparison of recognition of PrP in the normal and infected samples. The lanes containing unhydrolyzed S1 should contain relatively similar amounts of PrP protein recognized by the 3F4 antibody. The lanes containing the hydrolyzed, guanidine HCl-treated samples will allow the detection of PrP^{Sc} in the infected sample, since the only form left after the hydrolysis is the PrP^{Sc} form. The ratio of signal between the hydrolyzed and unhydrolyzed sample from the infected cow will determine the percentage of PrP that is in the PrP^{Sc} conformation. The hydrolyzed normal cow sample will further serve as a control that the hydrolysis of the PrP^C conformation was complete.

EXAMPLE 4

COMPARISON OF 3F4 AND R1 STAINING OF A SAMPLE

To determine the levels of PrP^{Sc} in affected mice, a prion infected mouse and a normal mouse are each sacrificed and their brains removed. A 10% (w/v) homogenate of each of the brains was prepared by dispersing the brain tissue in PBS. The brain homogenate was then subjected to a low speed centrifugation of 500 x g for 15 minutes to separate the suspended proteins from unwanted cellular debris. The total protein concentration of the supernatant (S1) was measured using a BCA Protein Assay (Pierce) and the concentration of each brain homogenate was adjusted with PBS to 3.5 mg/ml. A portion of the homogenate was saved to serve as a control of total brain proteins.

Each sample was then analyzed for PrP^{Sc} and PrP^C protein content by immunoblot. Two 100 μ l aliquot of each sample was processed and electrophoresed at 4°C on an 8% polyacrylamide slab gel as described in Example 2, i.e. under nondenaturing conditions. The gel is loaded with two lanes of sample from the affected mouse and two lanes of sample from

the control mouse, and preferable one lane of each is run on one side of the gel, one lane of each on the other side of the gel, with wells containing no sample separating the two sides. The gel is run and transferred to nylon as per Example 1.

Following transfer, the nylon is cut to separate the nylon into two separate blots, each containing a lane of affected and a lane of control sample. Each nylon filter is placed in a heat-sealable plastic bag, and 0.1 ml of blocking solution is added per square cm of filter. The blocking solution contains 5% (w/v) nonfat dry milk (Carnation) and 0.02% sodium azide in PBS. After 1 hour shaking at room temperature, antibody 3F4 is added to one blot in a 1:200 dilution, and antibody R1 is added to the other blot in a 1:200 dilution, and the filter is incubated for 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody are removed, and each filter washed three times.

The secondary antibody is properly labeled to allow immunoblot detection. The secondary antibody is added at a much more dilute level than the primary antibody, from 1:500 to 1:2000 dilution. After incubation, the filter is washed several times in PBS, each wash being about 10 minutes in length. The filter is then placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C.

The level of PrP^{Sc} in the affected sample can be determined by comparing the relative levels of signal of each antibody in the affected versus the control mouse. This process can be a physical comparison to determine the relative differences in signal level between the R1 and the 3F4 antibody, or it may be a quantitative comparison. Comparison techniques, both physical and quantitative, will be known to those skilled in the art. Quantitative comparison can be assessed using blot scanning techniques in which the levels are determined using computer programs specially designed to assess comparative levels. One such method is using a modified Excel spreadsheet program. Such programs allow for adjustments between samples based on variables such as background, time of hybridization, etc.

Levels of 3F4 staining should be consistent between the normal control samples of both the R1 and 3F4 antibody blots. The level of difference between R1 staining and 3F4 staining in the affected sample should allow the quantification of PrP^{Sc} in the sample by subtracting the level of PrP^C signal using the 3F4 antibody from the level of PrP^{Sc} and PrP^C signal from R1 antibody. Such quantification may be adjusted based on antibody sensitivity, concentration, etc.

EXAMPLE 5

DETECTION OF β A4 IN HUMAN AND MOUSE BRAIN

A number of mouse models for Alzheimer's disease exhibit many of the hallmark protein changes associated with the human disease. Two examples are: 1) mice with a modified human APP under the control of the PDGF promoter (the "Athena-Lilly mouse"),
5 the production and phenotype of these mice are described in U.S. Patent No.5,612,486, and 2) mice with a mutant isoform of human APP under the control of the prion gene promoter (the "Hsiao mouse") *see* Hsiao et al., *Science* 274:99-102 (1996). These mouse models for Alzheimer's disease display amyloid deposits and are capable of producing all three major
10 APP isoforms, and levels of APP and A β 40/A β 42 that progressively increase during the mouse's lifetime.

A 10% (w/v) brain homogenate from normal mice, affected Hsiao mice, and affected Athena mice are prepared in TBS (25 mM Tris). After a low speed centrifugation at 500 x g for 15 min, the total protein in the supernatant (S1) is measured using a BCA Protein Assay
15 (Pierce) and the concentration is adjusted to 2.5 mg/ml with PBS.

A Dispase digestion is done at an enzyme to protein ratio of 1:25. The protein is digested with 100 μ g/ml Dispase (Worthington) for 60 min at 37°C in the presence of 0.15% Zwittergent 3-12 (Calbiochem). The digestion is stopped by the addition of 50 mM EDTA. PAGE and transfer of the proteins is performed as in Example 1.

20 The nylon filter is placed in a heat-sealable plastic bag, and 0.1 ml of blocking solution is added per square cm of filter. The blocking solution contains 5% (w/v) nonfat dry milk (Carnation); 0.01% antifoam A; and 0.02% sodium azide in PBS. After 1 hour shaking at room temperature, a monoclonal antibody recognizing the β -amyloid protein, such as RDI-BAMYLOID (Research Daignostics) is added in a 1:200 dilution, and the filter incubated for
25 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody are removed, and the filter washed three times.

The filter is incubated with an anti-mouse Ig secondary antibody in blocking solution for 1-2 hours at room temperature. The secondary antibody is radiolabeled with to allow immunoblot detection. For the radiolabeled I¹²⁵ probe, approximately 10⁴ cm of the reagent is
30 added per square centimeter of filter. After incubation, the filter is washed several times in PBS, each wash being about 10 minutes in length. The filter is placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C.

The resulting blot will have a band corresponding to β A4 in the sample of the treated Athena Lilly or Hsiao mouse brain, both before and after digestion with Dispase. The sample of the normal mouse brain protein may show a low level band due to background with the APP protein. The lane with the normal brain sample subjected to hydrolysis with Dispase should not have a detectable band, since the Dispase will have hydrolyzed the normal protein in the sample.

This protocol may also be used with human brain or biopsy material, to identify and diagnose individuals suspected of having Alzheimer's disease.

10

EXAMPLE 6

DETECTION OF TTR IN BIOPSY MATERIAL

A biopsy sample from the liver is taken from an individual thought to be suffering from familial amyloid polyneuropathy (FAP). Homogenates (10% (w/v)) of the liver sample and a normal liver control sample are prepared in TBS (25 mM Tris). After a low speed centrifugation at 500 x g for 15 min, the total protein in the supernatant (S1) is measured using a BCA Protein Assay (Pierce) and the concentration is adjusted to 3.5 mg/ml with PBS.

A Neprilysin digestion is done at an enzyme to protein ratio of 1:35. The protein is digested with 100 μ g/ml Dispase for 60 min at 37°C in the presence of 0.2% Sarkosyl. The digestion is stopped by the addition of 50 mM EDTA. An aliquot of the proteins obtained in S1 both before and after the neprilysin digestion is electrophoresed at 4°C on an 8% polyacrylamide slab gel as described in Example 2, i.e. under nondenaturing conditions. The gel is run and proteins are transferred to nylon as in Example 1.

The nylon filter is placed in a heat-sealable plastic bag, and 0.1 ml of blocking solution is added per square cm of filter. The blocking solution contains 5% (w/v) nonfat dry milk (Carnation) and 0.02% sodium azide in PBS. After 1 hour shaking at room temperature, a monoclonal antibody recognizing the amyloid conformation of TTR is added in a 1:100 to 1:500 dilution, and the filter incubated for 2-4 hours at 4°C with gentle agitation on a platform shaker. Following incubation, the blocking solution and antibody are removed, and the filter washed three times.

The secondary antibody is enzyme-conjugated with horseradish peroxidase to allow immunoblot detection. The secondary antibody is added at a much more dilute level than the primary antibody, from 1:500 to 1:2000 dilution. After incubation, the filter is washed several

times in PBS, each wash being about 10 minutes in length. The filter is then placed in a cassette with a piece of Xomat X-ray film (Kodak) at -70°C .

The resulting blot will have a band corresponding to the amyloid conformation of TTR in an affected biopsy sample, but not in a biopsy sample in which the amyloids are not present. Moreover, the concentration of the TTR amyloid conformation may be a prognostic indicator as to the severity of the disease, and biopsies using multiple samples may also be used to determine disease progression.

The instant invention is shown and described herein in what is considered to be the most practical, and preferred embodiments. It is recognized, however, that departures may be made therefrom, which are within the scope of the invention, and that obvious modifications will occur to one skilled in the art upon reading this disclosure.

What is claimed is:

1. An assay method, comprising the steps of:
treating a sample suspected of containing a protein which assumes a first
5 conformation and a second, disease related conformation with a compound which hydrolyzes
the protein in the first conformation but not the second, disease related conformation to
provide a treated sample;
contacting the treating sample with a binding partner which binds the disease
related conformation; and
10 determining the presence of the disease related conformation based on binding of
the binding partner.
2. The assay method of claim 1, wherein the protein is a PrP protein, the
first conformation is PrP^C and the second, disease related conformation is PrP^{Sc} and the
15 compound which hydrolyzes the PrP^C changes the conformation of the PrP^{Sc} so that an
epitope not previously exposed is exposed after the treating.
3. The assay method of claim 2, wherein the compound which hydrolyzes
PrP^C but not PrP^{Sc} is selected from the group consisting of metalloendopeptidase and Dispace.
20
4. The assay method of claim 1, further comprising:
separating away protein of the second, disease related conformation from the
treated sample;
25 contacting the separated portion with the binding partner.
5. The assay method of claim 4, wherein the separating is carried out by
centrifuging and the binding partner is an antibody and wherein the protein is selected from
the group consisting of a PrP protein, a TTR protein and an amyloid protein.

30

6. The assay method of claim 1, further comprising:
pretreating the sample prior to treating the sample with the compound which
hydrolyzes the protein in the first conformation wherein the pretreating reduces the
concentration of protein in the sample other than the disease related conformation of the
5 protein.
7. The assay method of claim 6, wherein the pretreating comprises removing
protein in the sample by contacting the sample with antibodies bound to a support which
antibodies bind protein other than the protein in the second, disease related conformation.
10
8. The assay method of claim 6, wherein the pretreating comprises
hydrolyzing proteins in the sample other than proteins in the second, disease related
conformation.
9. The assay method of claim 1, wherein the binding partner is an R1
15 antibody.
10. An assay method, comprising the steps of :
treating a sample suspected of containing a protein which assumes a first
20 conformation and a second, disease related conformation with a compound which hydrolyzes
the protein in the first conformation but not the second, disease related conformation to
provide a treated sample;
denaturing protein in the second, disease related conformation to provide a
treated, denatured sample;
25 contacting the treated, denatured sample with a binding partner which binds the
denatured, second disease related conformation of the protein; and
detecting the second, disease related conformation of the protein based on
binding to the binding partner.
11. The assay method of claim 10, wherein the protein is a protein selected
30 from the group consisting of a PrP protein, a TTR protein and an amyloid protein.

12. The assay method of claim 10, wherein the protein is a PrP protein, the first conformation of the protein is PrP^C and the second, disease related conformation of the protein is PrP^{Sc}.

5 13. The assay method of claim 10, wherein the protein in the first conformation is PrP^C and the compound which hydrolyzes PrP^C is selected from the group consisting of Dispase and metalloendopeptidase.

10 14. The assay method of claim 10, wherein the binding partner is bound to a detectable label.

15 15. The assay method of claim 14, wherein the detecting is carried out using time-resolved disassociation enhanced fluorescence.

16. The assay method of claim 14, wherein the detecting is carried out using a dual wavelength, laser driven fluorometer.

17. An assay method, comprising the steps of:
providing a sample suspected of containing a protein which assumes a first
20 conformation and a second, disease related conformation;
contacting the sample with a first binding partner which first binding partner binds the first conformation but not the second, disease related conformation of the protein, wherein the first binding partner is added in sufficient amounts so as to bind to substantially all available epitopes in the sample;
25 contacting the sample with a second binding partner which binds to both the first conformation of the protein and the second, disease related conformation of the protein; and
detecting the presence of the second, disease related conformation of the protein based on binding to the second binding partner.

30 18. The assay of claim 17, wherein the first binding partner is a 3F4 antibody.

19. The assay method of claim 17, wherein the second binding partner is a R1 antibody.

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/US 99/23115

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 G01N33/68 G01N33/569 C07K14/47 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MEDLINE, Washington DC USA; abstract no. 94184819, abstract line 18 XP002129708 & W. GOLDMAN: " PrP gene and its association with spongiform encephalopathies." BRITISH MEDICAL BULLETIN, vol. 49, no. 4, 1 October 1993 (1993-10-01), pages 839-859, London UK	1-19
Y	WO 98 37210 A (KANTON ZÜRICH) 27 August 1998 (1998-08-27) cited in the application claims 1,2	1-19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 February 2000

Date of mailing of the international search report

17/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bohemen, C

INTERNATIONAL SEARCH REPORT

information on patent family members

Internal Application No

PCT/US 99/23115

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837210 A	27-08-1998	EP 0861900 A AU 6498698 A	02-09-1998 09-09-1998
